


518 Rec'd PCT/PTO 20 JUL 2001
09/889800

CERTIFICATION UNDER 37 CFR 1.10

I hereby certify that this Transmittal Letter and the papers indicated as being transmitted therewith are being deposited with the United States Postal Service on this date shown below in an envelope as "Express Mail Post Office to Addressee" under the below indicated Mailing Label Number, addressed to: Box PCT, Assistant Commissioner for Patents, Washington, D.C. 20231.

Mailing Label No.: EF232849690US

Deposit Date: July 20, 2001


Name: Janet Farr

ATTORNEY'S DOCKET No. DYOUP0219US

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
(DO/EO/US)**

In re national phase of:

Applicant(s): Richard Henry Williams et al.
International Application No.: PCT/GB00/00367
International Filing Date: 7 February 2000
Priority Date Claimed: 5 February 1999
Title of Invention: FERTILISER

**TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED
OFFICE (DO/EO/US) CONCERNING ENTRY INTO U.S. NATIONAL
PHASE UNDER 35 U.S.C. 371**

Box PCT
Assistant Commissioner for Patents
Washington D.C. 20231

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information under 35 U.S.C. 371:

1. This express request to immediately begin national examination procedures (35 U.S.C. 371(f)).
2. The U.S. National Fee (35 U.S.C. 371(c)(1)) and other fees (37 CFR 1.492) as indicated below.

3. A copy of the International application (35 U.S.C. 371(c)(2)):
- a. ☒ is transmitted herewith
(International Publication No. WO 00/46169).
 - b. ☐ is not required, as the application was filed with the United States Receiving Office.
 - c. ☐ has been transmitted by the International Bureau. A copy of Form PCT/IB/308 is enclosed.
4. ☐ A translation of the International application into the English language (35 U.S.C. 371(c)(2)) is transmitted herewith.
5. Amendments to the claims of the International application under PCT Article 19 (35 U.S.C. 371(c)(3)):
- a. ☒ are transmitted herewith.
 - b. ☐ have been transmitted by the International Bureau.
6. ☐ A translation of the amendments to the claims under PCT Article 19 (38 U.S.C. 371(c)(3)) is transmitted herewith.
7. A copy of the international examination report (PCT/IPEA/409)
- a. ☒ is transmitted herewith.
 - b. ☐ is not required as the United States Patent and Trademark Office was the IPEA.
8. Annex(es) to the international preliminary examination report
- a. ☒ is/are transmitted herewith.
 - b. ☐ is not required as the United States Patent and Trademark Office was the IPEA.
9. ☐ A translation of the annexes to the international preliminary examination report is transmitted herewith.
10. ☐ An oath or declaration of the inventor (35 U.S.C. 371(c)(4)) complying with 35 U.S.C. 115 is submitted herewith.

T022F0T"00222222

11. An International Search Report (PCT/ISA/210)
- a. ☒ is transmitted herewith.
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was searched by the United States International Searching Authority.
12. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98 is transmitted herewith, along with Form PTO-1449 and copies of citations listed.
13. ☐ An assignment document is transmitted herewith for recording, along with a separate cover sheet.
14. ☒ A preliminary amendment is enclosed.
15. ☐ A verified statement claiming small entity status is enclosed.
16. ☐ Other:

09/889800 101201

Basic National Fee					Fee
IPEA - US					\$690.00
ISA - US					\$710.00
PTO not ISA or IPEA					\$1,000.00
Claims meet PCT Art. 33(1)-(4) - IPEA - US					\$100.00
Filing with EPO or JPO search report					\$860.00
Enter appropriate basic fee →					\$860.00
Claims*	Number filed		Number extra	Rate	
Total claims	44	-20	22	\$18.00	\$432.00
Independent claims	3	-3	0	\$80.00	\$0.00
Multiple dependent claims (if applicable)				\$270.00	
Total of above					\$1,292.00
Small entity statement enclosed, 1 if Yes, 0 if No →				0	\$0.00
Total national fee					\$1,292.00
Fee for recording enclosed assignment				\$40.00	
Total fees enclosed					\$1,292.00

*After any attached preliminary amendment reducing the number of claims and/or deleting multiple dependencies.

☒ [X] A check in the amount of \$ 1,292.00 to cover the above fees is enclosed.

☐ [] Please charge our Deposit Account No. 18-0988 in the amount of \$ _____. A duplicate copy of this sheet is enclosed.

WARNING: TO AVOID ABANDONMENT OF THE APPLICATION THE BASIC NATIONAL FEE MUST BE PAID WITHIN THE 20/30 MONTH TIME LIMIT.

T02T01 0066860

16. The Commissioner is hereby authorized to charge the following additional fees that may be required by this paper and during the entire pendency of this application to our Deposit Account No. 18-0988:


a. ☒ 37 CFR 1.492(a)(1), (2), (3), (4) and (5) (filing fees)

WARNING: BECAUSE FAILURE TO PAY THE NATIONAL FEE WITHIN 30 MONTHS WITHOUT EXTENSION (37 CFR S 1.495(B)(2)) RESULTS IN ABANDONMENT OF THE APPLICATION, IT WOULD BE BEST TO ALWAYS CHECK THE ABOVE BOX.

b. ☐ 37 CFR 1.492(b), (c) and (d) (presentation of extra claims)

NOTE: Because additional fees for excess or multiple dependent claims not paid on filing or on later presentation must only be paid or these claims cancelled by amendment prior to the expiration of the time period set for response by the PTO in any notice of fee deficiency (37 CFR 1.492(d)), it might be best not to authorize the PTO to charge additional claim fees, except possibly when dealing with amendments after final action.

Respectfully submitted,



Neil A. DuChez
Reg. No. 26,725

Direct all correspondence and telephone calls to:

Neil A. DuChez, Esq.
RENNER, OTTO, BOISSELLE & SKLAR, LLP
1621 Euclid Avenue, 19th Floor
Cleveland, Ohio 44115
Tel: 216-621-1113 Fax: 216-621-6165

09/889800 101201

ATTORNEY'S DOCKET No. DYOUP0219US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE
(DO/EO/US)

In re national phase of:

Applicant(s): Richard Henry Williams et al.
International Application No.: PCT/GB00/00367
International Filing Date: 7 February 2000
Priority Date Claimed: 5 February 1999
Title of Invention: FERTILISER

PRELIMINARY AMENDMENT

Box PCT
Assistant Commissioner for Patents
Washington D.C. 20231

Sir:

Prior to commencing examination of the above-identified application, kindly enter the following amendment.

In the Claims:

Kindly amend claims 7-11, 13, 14, 17, 19, 21 and 22 as follows:

7. (Amended) A fertiliser composition according to claim 1 wherein the phosphonate is ammonium, sodium or potassium phosphonate or a mixture thereof.

8. (Amended) A fertiliser composition according to claim 1 wherein the thiosulphate is ammonium, sodium or potassium thiosulphate or a mixture thereof.

9. (Amended) A fertiliser composition according to claim 2 wherein the derivative of salicylic acid is salicylamide or a salt thereof.

10. (Amended) A fertiliser composition according to claim 2 wherein the homologue of salicylic acid is benzoic acid or a salt or derivative thereof.

09/889800 JC17 Rec'd PCT/PTO 20 JUL 2001

11. (Amended) A fertiliser composition according to claim 2 wherein the salt of salicylic acid, its homologue or derivative is an organic or inorganic salt.

13. (Amended) A fertiliser composition according to claim 1 in the form of a concentrate.

14. (Amended) A fertiliser composition according to claim 1 in the form of an aqueous solution.

17. (Amended) A fertiliser composition according to claim 1 further comprising a plant growth regulator.

19. (Amended) A method for fertilising a plant comprising applying a fertiliser composition according to claim 1 to the plant or its environs.

21. (Amended) A method according to claim 19 wherein the thiosulphate is applied at 250 g/ha to 6 kg/ha.

22. (Amended) A method according to claim 19 wherein at least one salicylic acid, a homologue, derivative, or salt thereof is applied at 1 g/ha to 100 g/ha.

A version of the above-amended claims marked to indicate the specific amendments may be found in the attached Appendix, in accordance with 37 C.F.R. 1.121(c)(1).

Kindly cancel claims 23 and 24, without prejudice.

Kindly add claims 25-46 as follows:

25. A fertiliser composition according to claim 3 wherein the phosphonate is ammonium, sodium or potassium phosphonate or a mixture thereof.

26. A fertiliser composition according to claim 3 wherein the derivative of salicylic acid is salicylamide or a salt thereof.

27. A fertiliser composition according to claim 3 wherein the homologue of salicylic acid is benzoic acid or a salt or derivative thereof.

28. A fertiliser composition according to claim 3 wherein the salt of salicylic acid, its homologue or derivative is an organic or inorganic salt.

29. A fertiliser composition according to claim 3 in the form of a concentrate.

30. A fertiliser composition according to claim 3 in the form of an aqueous solution.

31. A fertiliser composition according to claim 3 further comprising a plant growth regulator.

32. A method for fertilising a plant comprising applying a fertiliser composition according to claim 3 to the plant or its environs.

33. A method according to claim 32 wherein at least one salicylic acid, a homologue, derivative, or salt thereof is applied at 1 g/ha to 100 g/ha.

34. A fertiliser composition according to claim 6 wherein the phosphonate is ammonium, sodium or potassium phosphonate or a mixture thereof.

35. A fertiliser composition according to claim 5 wherein the thiosulphate is ammonium, sodium or potassium thiosulphate or a mixture thereof.

36. A fertiliser composition according to claim 5 wherein the derivative of salicylic acid is salicylamide or a salt thereof.

37. A fertiliser composition according to claim 5 wherein the homologue of salicylic acid is benzoic acid or a salt or derivative thereof.

38. A fertiliser composition according to claim 5 wherein the salt of salicylic acid, its homologue or derivative is an organic or inorganic salt.

39. A fertiliser composition according to claim 5 in the form of a concentrate.

40. A fertiliser composition according to claim 5 in the form of an aqueous solution.

41. A fertiliser composition according to claim 5 further comprising a plant growth regulator.

42. A method for fertilising a plant comprising applying a fertiliser composition according to claim 5 to the plant or its environs.

43. A method according to claim 42 wherein the thiosulphate is applied at 250 g/ha to 6 kg/ha.

44. A method according to claim 42 wherein at least one salicylic acid, a homologue, derivative, or salt thereof is applied at 1 g/ha to 100 g/ha.

45. A method of stimulating the growth of a plant comprising applying an effective amount of the fertiliser composition of claim 1 to the plant or its environs to stimulate the growth of the plant.

46. A method of controlling parasitic fungi comprising applying the fertiliser composition of claim 1 to the parasitic fungi.


REMARKS

Claims 7-11, 13, 14, 17, 19, 21 and 22 have been amended. Claims 23 and 24 have been canceled, without prejudice. Claims 25-46 have been added. Claims 1-22 and 25-46 are in the application upon entry of this amendment. Entry of this amendment is respectfully requested.

Claims 7-11, 13, 14, 17, 19, 21 and 22 have been amended to convert the claims from multiple dependent claims to singularly dependent claims. Claims 25-44 have been added to provide for claim coverage using singular dependant claim format rather than multiple dependent claim format as originally presented. This is being done to reduce the government filing fee.

Claims 23 and 24 have been canceled and recast as claims 45 and 46, respectively. This has been done to convert the claims to a form that is acceptable to USPTO requirements.

Respectfully submitted,



Neil A. DuChez
Reg. No. 26,725

Direct all correspondence and telephone calls to:

Neil A. DuChez, Esq.
RENNER, OTTO, BOISSELLE & SKLAR, LLP
1621 Euclid Avenue, 19th Floor
Cleveland, Ohio 44115
Tel: 216-621-1113 Fax: 216-621-6165

In the Claims:

7. (Amended) A fertiliser composition according to [any one of claims 1-4 and 6] claim 1 wherein the phosphonate is ammonium, sodium or potassium phosphonate or a mixture thereof.

8. (Amended) A fertiliser composition according to [any one of claims 1 and 4-7] claim 1 wherein the thiosulphate is ammonium, sodium or potassium thiosulphate or a mixture thereof.

9. (Amended) A fertiliser composition according to [any one of claims 2-8] claim
2 wherein the derivative of salicylic acid is salicylamide or a salt thereof.

10. (Amended) A fertiliser composition according to [any one of claims 2-8] claim
2 wherein the homologue of salicylic acid is benzoic acid or a salt or derivative thereof.

11. (Amended) A fertiliser composition according to [any of claims 2-9] claim 2 wherein the salt of salicylic acid, its homologue or derivative is an organic or inorganic salt.

13. (Amended) A fertiliser composition according to [any preceding] claim 1 in the form of a concentrate.

14. (Amended) A fertiliser composition according to [any one of claims 1-12] claim 1 in the form of an aqueous solution.

17. (Amended) A fertiliser composition according to [any preceding] claim 1 further comprising a plant growth regulator.

19. (Amended) A method for fertilising a plant comprising applying a fertiliser composition according to [any preceding] claim 1 to the plant or its environs.

21. (Amended) A method according to claim 19 [or claim 20] wherein the thiosulphate is applied at 250 g/ha to 6 kg/ha.

22. (Amended) A method according to [any one of claims 19 to 21] claim 19 wherein at least one salicylic acid, a homologue, derivative, or salt thereof is applied at 1 g/ha to 100 g/ha.

Claims 23 and 24 have been cancelled.

Claims 25-46 have been added.

Fertiliser

The present invention relates to novel compositions having fertilising and anti-fungal effects, to processes for their preparation, and to methods of fertilising plants and controlling fungi using them.

5 Phosphorus is one of the essential major elements required by plants and it is usually supplied to plants in the form of phosphate and/or polyphosphate. Phosphates are the salts of phosphoric acid (having the formula H_3PO_4 and molecular weight of 98). In recent years, it has been shown that plants can obtain phosphorus from phosphonates (sometimes also referred to as phosphites) which
10 are the salts (organic or inorganic) of phosphonic acid (also referred to as phosphorous acid) (having the formula H_3PO_3 and molecular weight of 82). See, for example, US Patent Nos. 5,514,200 & 5,830,255 to Lovatt; US Patent No. 5,707,418 to Hsu; US Patent No. 5,800,837 to Taylor. These describe formulations containing phosphorous acid or phosphonates suitable as fertilisers
15 for plants. It has also been shown that phosphonate compounds are useful as fungicides, especially where the fungal organisms are phycomycetes or oomycetes. See, for example, US Patent Nos. 4,075,324 & 4,119,724 to Thizy; US Patent No. 4,139,616 to Lacroix et al; US Patent Nos. 4,698,334, 4,806,445 & 5,169,646 to Horriere et al; US Patent Nos 4,935,410 & 5,070,083 to Bartlet; US
20 Patent No. 5,736,164 to Taylor. These describe formulations, containing phosphorous acid or phosphonates, suitable as fungicides for plants.

Ammonium thiosulphate and potassium thiosulphate, either alone or mixed with other liquid fertiliser components, have been used for many years as fertilisers. See literature on "Thio-sul"® and KTS® sulphur fertilisers produced
25 by Tessenderlo Kerley. See also UK Patent No. GB 2,259,912 to Sampson, which describes the use of ammonium thiosulphate in a plant growth stimulator.

WPI Abstract Accession No. 91-249421 discloses a cut flower preserving agent comprising a water-soluble silver salt (100 pts. wt.) and thiosulphate (300-2500 pts. wt.) to which a phosphite (30-300 pts. wt.) is added as a stabilising
30 agent.

09889800-101201

Some of the problems with the prior art are that the fertilising effect of phosphonate is less than might be expected from the amount of phosphorus applied, and the fungicidal effect is fairly limited in terms of the types of pathogen controlled. This is due to a complex mode of action involving a combination of some fungistatic action and natural plant defences coming into play (See Guest D I & Grant B R (1991) - The Complex action of phosphonates in plants - Biological Reviews 66, 159-187). The use of phosphonate, whilst improving the resistance of plants to infections of downy mildew (eg Plasmopora) and Phytophthora diseases, does tend to increase the risk of ascomycete (eg Erysiphe) infections. The present invention seeks to provide a solution to these problems.

According to one aspect of the present invention there is provided a fertiliser composition comprising at least one phosphonate and at least one thiosulphate.

According to another aspect of the present invention there is provided a fertiliser composition comprising at least one phosphonate and at least one salicylic acid, homologue, derivative, or salt thereof.

According to yet another aspect of the present invention there is provided a fertiliser composition comprising at least one thiosulphate and at least one salicylic acid, homologue, derivative, or salt thereof.

According to a further aspect of the present invention there is provided a fertiliser composition comprising at least one thiosulphate, at least one phosphonate and at least one salicylic acid, homologue, derivative, or salt thereof.

The present invention comprises using a mixture of a phosphonate together with either a thiosulphate, or at least one salicylic acid, homologue, derivative, or salt thereof. The use of this combination shows a synergistic effect, in that the combination of phosphonate with thiosulphate or salicylic acid, homologues, salts or derivatives thereof produces a greater fertiliser effect and fungicidal effect than the individual components used separately. There may be

3

an even greater effect if all three components (ie phosphonate, thiosulphate and salicylic acid, homologue, salt or derivative thereof) were used together. The combination of thiosulphate with a salicylic acid, homologue, salt or derivative thereof, in the absence of phosphonate, also produces a fertiliser effect and fungicidal effect.

Fertilisers based on the present invention provide a greater growth effective response than phosphonates or thiosulphates alone and the degree of fungicidal protection or resistance is broader than that achieved with phosphonates or thiosulphates alone. Plants treated with the present invention suffer less from phycomycete diseases (for example *phytophthoras* and downy mildews) than those treated with for example phosphonate alone and are also less prone to other parasitic fungi such as powdery mildews. Thus the present invention provides a means for applying a single product to plants which is an effective fungicide as well as an effective fertiliser.

Another advantage of the present invention is that the formulation is very storage stable, for example tests on mixtures of potassium phosphonate and ammonium thiosulphate stored for over one year have shown that there is no oxidation of the phosphonate to phosphate and the stored material shows no signs of cloudiness or precipitation. The use of further organic acids as buffers (as is required in US Patent Nos. 5,514,200 & 5,830,255) is also not required to achieve stable solutions.

By "phosphonate" we mean a salt of phosphonic acid (H_3PO_3). Phosphonates contain the trivalent $\equiv\text{PO}_3$ radical. For the avoidance of doubt, phosphonic acid is sometimes referred to as phosphorous acid and its salts as phosphites. Mixtures of phosphonates may be employed.

The phosphonate may be any metal ion or other cation which forms such a salt. As phosphonic acid has a P-H bond it forms a mono and di series of salts. Both mono and di salts and mixtures thereof may be used in the present invention. Preferably the phosphonate is an ammonium phosphonate or alkali phosphonate. Amongst the alkali phosphonates, sodium or potassium

phosphonate are preferred. Potassium phosphonate is particularly preferred, in the form of mono- and/or di-potassium phosphonate (KH_2PO_3 , K_2HPO_3 respectively).

Phosphonates may be produced by the neutralisation of phosphonic acid by an alkali. The present invention also encompasses the use of phosphonic acid which is subsequently converted to its phosphonate; this conversion may take place in situ or ex situ. When using, for example, potassium hydroxide for the neutralisation, depending on the molar ratio of potassium hydroxide to phosphorous acid, the phosphonate solution will contain varied amounts of di-potassium phosphonate, mono-potassium phosphonate and un-reacted phosphorous acid. We have found that an approximately 42% w/w solution, having a pH of between 6.7 and 7.3 and containing approximately equal amounts of mono- and di-potassium phosphonate is a clear, colourless and very stable starting material for our present invention.

The thiosulphate may be any suitable salt of a metal or other cation. Preferably the thiosulphate is ammonium, sodium or potassium thiosulphate or a mixture thereof. More preferably the thiosulphate is in the form of either ammonium or potassium thiosulphate ($(\text{NH}_4)_2\text{S}_2\text{O}_3$ or $\text{K}_2\text{S}_2\text{O}_3$).

The most common form of thiosulphate is ammonium thiosulphate, and this is readily available commercially as a 60% w/w solution, with a pH of about 7.5 and a specific gravity of about 1.32. If a higher proportion of potassium is required in the final foliar fertiliser, the ammonium thiosulphate can be substituted, either partly or wholly, with potassium thiosulphate.

The present invention includes functional homologues and derivatives of salicylic acid and its salts. By this we mean that the functional homologue or derivative should be capable of providing a fertiliser effect and/or antifungal effect. Examples of such derivatives of salicylic acid include salicylamide or a salt thereof, and esters.

Examples of homologues of salicylic acid include benzoic acid or a salt or derivative thereof, such as an ester. Examples of benzoic acid compounds which may be used in the present invention may be found in WO99/25191.

The salicylic acid is preferably in the form of its potassium salicylate salt or salicylamide - $C_7H_5KO_3$ or $C_7H_7NO_2$.

Salicylic acid itself has low solubility, but inorganic salts of salicylic acid, such as sodium or potassium salicylate are readily soluble. When salicylamide is used, rather than salicylic acid or a salicylate, the addition of a few drops of alkali assists in its solution, by forming for example sodium or potassium salicylamide. Salicylamide also dissolves more readily in the thiosulphate solution, the presence of small amounts of alkali or ammonia in the thiosulphate solution assisting in the solubilisation.

The preparation of the compounds used in the present invention is well known in the art. The compounds may be prepared in situ or ex situ.

In one embodiment, the composition of the present invention does not include a water-soluble silver salt. In another embodiment, if the composition contains a solution of 100 parts by weight water-soluble silver salt, and 300-2500 parts by weight thiosulphate, then the amount of phosphonate is other than 30 to 300 parts by weight

The compositions of the present invention are useful as fertiliser, particularly foliar fertilisers. More particularly the compositions of the present invention increase plant growth compared to the individual components alone, stimulate growth in plants, plant vigour and/or effect crop yield, for example by reducing tuber blight.

The compositions of the present invention also have an antifungal effect. This may be a fungicidal or fungistatic effect. The compositions of the present invention may have activity against parasitic fungi. The compositions may have activity against phycomycete diseases such as *phytophthoras* and downy mildews, for example, *Plasmopora*; and/or ascomycetes such as, for example, *Erysiphe*.

6

In one particularly preferred embodiment the composition further comprises a plant growth regulator. Preferably the plant growth regulator is chlormequat.

In order to apply the composition to the plant or environs of the plant, the composition may be used as a concentrate or more usually is formulated into a composition which includes an effective amount of the composition of the present invention together with a suitable inert diluent, carrier material and/or surface active agent. Preferably the composition is in the form of an aqueous solution which may be prepared from the concentrate. By effective amount we mean that the composition (and/or its individual components) provides a fertilising and/or antifungal effect. Preferably an effective amount of the components is a concentration of up to about 4M phosphonate, up to about 5M thiosulphate and/or up to about 0.8M salicylate. Thus, in one embodiment the concentrate may comprise up to about 10M of the components. The concentrate formulation may for example be diluted at ratios of concentrate to water of about 1:40 to 1:600, and generally is formulated to have pH of about 6.5 to 8.5. At a 1:40 dilution, a concentrate of about 10M would give rise to an application concentrate of up to about 0.25M.

The rate and timing of application will depend on a number of factors known to those skilled in the art, such as the type of species etc.

The composition is generally applied in an amount of from 0.01 to 10kg per hectare, preferably 0.1 to 6kg per hectare. Preferably the phosphonate is applied at 150 g/ha to 2 kg/ha. Preferably the thiosulphate is applied at 250 g/ha to 6 kg/ha. Preferably the salicylic acid, a homologue, derivative, or salt thereof is applied at 1 g/ha to 100 g/ha.

In one preferred embodiment, a fertiliser composition according to the present invention comprises about 150 g/l phosphonate, about 275 g/l thiosulphate and/or about 10 g/l salicylamide. Preferably the phosphonate comprises about 75 g/l mono-potassium phosphonate and about 75 g/l di-potassium phosphonate.

As well as varying amounts of each compound to be blended together, as is common with many foliar fertilisers, it is also possible to combine other fertilising elements, such as but not limited to, iron, copper, boron and molybdenum (often known as micronutrients) in the final solution. These may be added as soluble inorganic compounds (eg sodium borate or sodium molybdate) or as chelates (eg copper EDTA) or other metal complexes.

The compositions of the present invention can be applied to the soil, plant, seed, or other area to be protected. Preferably the present invention is applied to the foliage of plants. The composition may be applied in the form of dusting powders, wettable powders, granules (slow or fast release), emulsion or suspension concentrates, liquid solutions, emulsions, seed dressings, or controlled release formulations such as microencapsulated granules or suspensions, soil drench, irrigation component, or preferably a foliar spray.

Dusting powders are formulated by mixing the active ingredient with one or more finely divided solid carriers and/or diluents, for example natural clays, kaolin, pyrophyllite, bentonite, alumina, montmorillonite, kieselguhr, chalk, diatomaceous earths, calcium phosphates, calcium and magnesium carbonates, sulfur, lime, flours, talc and other organic and inorganic solid carriers.

Granules are formed either by absorbing the active ingredient in a porous granular material for example pumice, attapulgite clays, fuller's earth, kieselguhr, diatomaceous earths, ground corn cobs, and the like, or on to hard core materials such as sands, silicates, mineral carbonates, sulfates, phosphates, or the like. Agents which are commonly used to aid in impregnation, binding or coating the solid carriers include aliphatic and aromatic petroleum solvents, alcohols, polyvinyl acetates, polyvinyl alcohols, ethers, ketones, esters, dextrans, sugars and vegetable oils, with the active ingredient. Other additives may also be included, such as emulsifying agents, wetting agents or dispersing agents.

Microencapsulated formulations (microcapsule suspensions CS) or other controlled release formulations may also be used, particularly for slow release over a period of time, and for seed treatment.

Alternatively the compositions may be in the form of liquid preparations to be used as dips, irrigation additives or sprays, which are generally aqueous dispersions or emulsions of the active ingredient in the presence of one or more known wetting agents, dispersing agents or emulsifying agents (surface active agents). The compositions which are to be used in the form of aqueous dispersions or emulsions are generally supplied in the form of an emulsifiable concentrate (EC) or a suspension concentrate (SC) containing a high proportion of the active ingredient or ingredients. An EC is an homogeneous liquid composition, usually containing the active ingredient dissolved in a substantially non-volatile organic solvent. An SC is a fine particle size dispersion of solid active ingredient in water. To apply the concentrates they are diluted in water and are usually applied by means of a spray to the area to be treated.

Suitable liquid solvents for ECs include methyl ketone, methyl isobutyl ketone, cyclohexanone, xylenes, toluene, chlorobenzene, paraffins, kerosene, white oil, alcohols (for example, butanol), methylnaphthalene, trimethylbenzene, trichloroethylene, N-methyl-2-pyrrolidone and tetrahydrofurfuryl alcohol (THFA).

These concentrates are often required to withstand storage for prolonged periods and after such storage, to be capable of dilution with water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. The concentrates may contain 1-85% by weight of the active ingredient or ingredients. When diluted to form aqueous preparations such preparations may contain varying amounts of the active ingredient depending upon the purpose for which they are to be used.

The composition may also be formulated as powders (dry seed treatment DS or water dispersible powder WS) or liquids (flowable concentrate FS, liquid seed treatment LS), or microcapsule suspensions CS for use in seed treatments. The formulations can be applied to the seed by standard techniques and through conventional seed treaters. In use the compositions are applied to the plants, to

the locus of the plants, by any of the known means of applying fertiliser compositions, for example, by dusting, spraying, or incorporation of granules.

When the final solution is to be applied to plants which, because of their hairy or waxy surface, may be difficult to wet, it may also be advantageous to include other additives, commonly known in the agrochemical industry, such as surfactants, wetting agents, spreaders and stickers. (Examples of wetting agents include silicone surfactants, nonionic surfactants such as alkyl ethoxylates, anionic surfactants such as phosphate ester salts and amphoteric or cationic surfactants such as fatty acid amido alkyl betaines).

As indicated above, the compounds of the invention may be the sole active ingredient of the composition or they may be admixed with one or more additional active ingredients such as nematicides, insecticides, synergists, herbicides, additional fungicides, additional fertilisers or plant growth regulators where appropriate.

As indicated above, the fertilisers produced according to this present invention are usually applied to the foliage of plants but may also be applied to the soil or added to the irrigation water. The fertilisers may be used advantageously on many types of agricultural and horticultural crops, including but not limited to, cereals, legumes, brassicas, cucurbits, root vegetables, sugar beet, grapes, citrus & other fruit trees and soft fruits. More particularly, crops that will benefit from the fertiliser include, but are not limited to, peas, oil seed rape, carrots, spring barley, avocado, citrus, mango, coffee, deciduous tree crops, grapes, strawberries and other berry crops, soybean, broad beans and other commercial beans, corn, tomato, cucurbitis and other cucumis species, lettuce, potato, sugar beets, peppers, sugar cane, hops, tobacco, pineapple, coconut palm and other commercial and ornamental palms, rubber and other ornamental plants.

Various further preferred features and embodiments of the invention will now be described by reference to the following non-limited Examples.

Example 1

Solution 1

An aqueous solution containing a total of 30% by weight of mono and di-
5 potassium phosphonate in roughly equal proportions.

Solution 2

An aqueous solution containing 55% by weight of ammonium thiosulphate ("ATS").

Solution 3

10 An aqueous solution containing 20 grams per litre of potassium salicylamide.

Solution 4

An aqueous solution containing 75 g/L mono potassium phosphonate, 75 g/L di-potassium phosphonate, 275 g/L ammonium thiosulphate and 10 g/L potassium salicylamide.

These solutions were applied to lettuce plants, both alone and in combination, and the applications were repeated after a 10 day interval. There were five replicates of each treatment and the results are presented as means of the five replicates. Five plants were also left unsprayed as an untreated control to the other treatments. After eight, twelve, sixteen and twenty-one days, the plants were examined for disease.

Table 1. Powdery Mildew Score (0 - 9, where higher number equals greater degree of disease)

Treatment (Applied initially and repeated 10 days later)	Days after first spray		
	8 days	12 days	16 days
Untreated	4.0	6.6	7.2
Solution 1 (1L/ha)	0.8	2.2	3.6
Solution 2 (1L/ha)	1.0	1.2	2.4
Solution 3 (1L/ha)	1.6	3.2	4.2
Solution 1 (1L/ha) + Solution 2 (1L/ha)	0.0	0.4	1.0
Solution 1 (1L/ha) + Solution 3 (1L/ha)	0.4	0.6	1.0
Solution 1 (1L/ha) + Solution 2 (1L/ha) + Solution 3 (1L/ha)	0.8	0.6	0.6

- 5 Table 1 shows the synergistic effect on disease levels achieved by adding Solutions 1 & 2 (phosphonate + ATS) and between Solutions 1 & 3 (phosphonate + salicylamide) and the further effect of using all three solutions together. Disease levels were reduced from a mean of 7.2 to a mean of 0.6

- 10 As well as assessing disease levels, the growth of the plants was assessed by measuring the mean plant diameters after 35 days growth and by measuring the mean above ground fresh and dry weights.

Table 2. Plant Growth after treatment with the example solutions

Treatment (Applied initially and repeated 10 days later)	Amount of Rooting (0-9, 0= least rooting) – mean	Plant Diameter (mm) -mean	Above-Ground Fresh Weight (g) – mean	Above-Ground Dry Weight (g) – mean
Untreated	5.3	124	102.3	8.3
Solution 1 (1L/ha)	6.0	148	116.3	9.3
Solution 2 (1L/ha)	5.3	160	109.0	8.7
Solution 3 (1L/ha)	4.7	150	104.7	8.5
Solution 1 (1L/ha) + Solution 2 (1L/ha)	6.7	144	119.0	9.5
Solution 1 (1L/ha) + Solution 3 (1L/ha)	6.7	170	120.7	9.7
Solution 1 (1L/ha) + Solution 2 (1L/ha) + Solution 3 (1L/ha)	6.7	168	131.7	10.6

- 5 Table 2 shows the synergistic effect on plant growth caused by adding Solutions 1 & 2 (phosphonate + ATS), Solutions 1 & 3 (phosphonate + salicylamide) and the further effect of using all three solutions together.

The abbreviations used in the following Examples A-E are:

A = phosphonate + thiosulphate

B = phosphonate + salicylate/salicylamide

5 C = thiosulphate + salicylate/salicylamide

D = thiosulphate + salicylate/salicylamide + phosphite

E = thiosulphate + salicylate + chlormequat

KP40 = 40% potassium phosphonate

10 KT47 = 47% potassium thiosulphate (w/v)

KS20 = 20% potassium salicylate (w/v)

CS8 = salicylamide (20g/l)

AT60 = 60% ammonium thiosulphate

PF723 = 55% ammonium thiosulphate

15

Examples A

Solution 1 = KP40 at 0.75 l/ha every 10 days

Solution 2 = PF723 at 1.0 l/ha every 10 days

20

Table A1

Percent Powdery Mildew - Lettuce

Treatment (Applied initially and repeated after a 10 day interval)	Days After First Spray		
	8 Days	12 Days	16 Days
Untreated	40	66	72
Solution 1 (1L/Ha)	8	22	36
Solution 2 (1L/Ha)	10	12	24
Solution 1 (1L/Ha) + Solution 2 (1L/Ha)	0	4	10

Table A2**Percent Powdery Mildew and Fertiliser Attributes – Sugar Beet**

Treatment (Applied initially and repeated after a 10 day interval)	Days After First Spray		Above Ground Fresh Weight (g) - Mean (x 1 Rate)
	+ 28 Days (x 1 Rate)	+ 35 Days (x 1 Rate)	
Untreated	22	31	144.7
Solution 1	6	4	152.3
Solution 2	2	2	153.3
Solution 1 + Solution 2	0	0	154.3

Table A3**Fertiliser Attributes – Spring Barley**

Treatment (Applied initially and repeated after a 10 day interval)	Plant Health 'Greenness' Score (0-9) +35 Days (x 1 Rate)	Above Ground Fresh Weight (g) - Mean (x 1 Rate)	Above Ground Dry Weight (g) - Mean (x 1 Rate)	Amount of Above Ground Tissue (0-9) at Harvest (x 1 Rate)
Untreated	5.6	68.8	7.7	5.0
Solution1	5.6	67.0	7.4	5.3
Solution 2	5.4	68.3	7.2	5.3
Solution 1 + Solution 2	6.6	72.7	7.9	6.0

Table A4

GRAPE TRIAL
Percent Powdery Mildew

Treatment (Applied initially and repeated at 10 day intervals)	Days After First Spray					
	+ 16 Days	+ 20 Days	+ 24 Days	+ 28 Days	+ 32 Days	+ 36 Days
Untreated	7.3	9.3	11.3	14.0	14.0	14.7
KP40 @ 0.75 l/ha	5.3	6.7	8.0	11.3	10.7	10.7
PF723 @ 1.0 l/ha	2.7	7.3	7.3	8.0	6.7	6.7
KP40 @ 0.75 l/ha + PF723 @ 1.0 l/ha	1.3	5.3	6.0	6.0	5.3	5.3

Table A5

LETTUCE TRIAL
Percent Powdery Mildew

Treatment (Applied initially and repeated at 10 day intervals)	Days After First Spray		
	+ 16 Days	+ 20 Days	+ 24 Days
Untreated	10.0	16.7	24.7
KP40 @ 0.75 l/ha	7.3	12.0	16.0
PF723 @ 1.0 l/ha	8.0	12.0	15.3
KP40 @ 0.75 l/ha + PF723 @ 1.0 l/ha	5.3	8.7	14.0

Table A6

Broad Bean – Fertiliser Attributes

(KP40 = 40% Potassium Phosphite; PF723 = 55% Ammonium Thiosulphate)

Treatment (2 Applications in total - every 15 days)	Above Ground Fresh Weight (g) - Mean	Above Ground Dry Weight (g) - Mean
Untreated	143.8	14.6
KP40 (5.0 l/ha) + PF723 (0.5 l/ha)	150.7	15.5
KP40 (3.0 l/ha) + PF723 (0.5 l/ha)	160.3	16.1
KP40 (5.0 l/ha) + PF723 (1.0 l/ha)	168.0	17.3
KP40 (3.0 l/ha) + PF723 (1.0 l/ha)	161.3	16.8

T02T01" 00868860

Table A7

Sugar Beet

Percent Powdery Mildew and Fertiliser Attributes

(KP40 = 40% Potassium Phosphite; AT60 = 60% Ammonium Thiosulphate)

Treatment (2 Applications in total - every 15 days)	Days After First Spray		Root Fresh Weight (g) - Mean	Root Dry Weight (g) - Mean	Above Ground Fresh Weight (g) - Mean	Above Ground Dry Weight (g) - Mean
	+ 24 Days	+ 28 Days				
Untreated	14	24	83.0	12.2	137.8	15.3
KP40 (0.375 l/ha) + AT60 (10.0 l/ha)	0	4	90.3	12.9	142.3	15.3
KP40 (0.75 l/ha) + AT60 (10.0 l/ha)	0	2	91.7	13.9	145.0	16.4
KP40 (0.375 l/ha) + AT60 (6.0 l/ha)	0	0	82.7	12.4	144.0	16.5
KP40 (0.75 l/ha) + AT60 (6.0 l/ha)	0	0	97.7	13.6	145.3	15.5
KP40 (3.75 l/ha) + AT60 (10.0 l/ha)	0	6	89.7	13.2	144.0	15.8
KP40 (2.5 l/ha) + AT60 (10.0 l/ha)	2	8	89.7	14.0	143.3	15.4
KP40 (2.5 l/ha) + AT60 (6.0 l/ha)	0	4	98.0	13.9	146.7	16.1

Table A8

Sugar Beet

Percent Powdery Mildew and Fertiliser Attributes

(KP40 = 40% Potassium Phosphate; AT60 = 60% Ammonium Thiosulphate)

Treatment (2 Applications in total - every 15 days)	Days After First Spray		Root Fresh Weight (g) - Mean	Root Dry Weight (g) - Mean	Above Ground Fresh Weight (g) - Mean	Above Ground Dry Weight (g) - Mean
	+ 24 Days	+ 28 Days				
Untreated	22	28	109.7	13.3	144.7	14.7
KP40 (0.375 l/ha) + AT60 (10.0 l/ha)	0	2	110.3	13.2	149.7	15.5
KP40 (0.75 l/ha) + AT60 (10.0 l/ha)	0	2	118.7	13.8	147.7	15.1
KP40 (0.375 l/ha) + AT60 (6.0 l/ha)	0	6	117.7	13.6	151.3	15.0
KP40 (0.75 l/ha) + AT60 (6.0 l/ha)	2	2	113.7	13.7	150.7	15.2
KP40 (3.75 l/ha) + AT60 (10.0 l/ha)	0	0	119.0	14.2	150.7	15.0
KP40 (2.5 l/ha) + AT60 (10.0 l/ha)	2	0	117.3	14.7	148.7	15.0
KP40 (2.5 l/ha) + AT60 (6.0 l/ha)	2	2	119.7	14.0	154.3	15.6

Examples B

Solution 1 = KP40 at 0.75 l/ha

Solution 3 = CS8 at 1.0 l/ha

5

Table B1

Fertiliser Attributes – Strawberry

Treatment (Applied initially and repeated after a 10 day interval)	Plant Health 'Greenness' Score (0-9) + 28 Days (x 1 Rate)	Above Ground Fresh Weight (g) - Mean (x 1 Rate)	Above Ground Dry Weight (g) - Mean (x 1 Rate)
Untreated	4.8	50.8	5.0
Solution 1	5.6	53.7	5.1
Solution 3	6.0	56.0	5.6
Solution 1 + Solution 3	6.4	63.0	6.2

Table B2

Fertiliser Attributes – Spring Barley

Treatment (Applied initially and repeated after a 10 day interval)	Above Ground Fresh Weight (g) - Mean (x 1 Rate)	Above Ground Dry Weight (g) - Mean (x 1 Rate)	Amount of Above Ground Tissue (0-9) (x 1 Rate)
Untreated	77.3	8.6	5.0
Solution 1	78.3	8.7	5.3
Solution 3	75.0	8.4	5.3
Solution 1 + Solution 3	81.7	9.7	5.7

Table B3

Percent Powdery Mildew – Sugar Beet

Treatment (Applied initially and repeated after a 10 day interval)	Days After First Spray	
	+ 28 Days (x 1 Rate)	+ 32 Days (x 1 Rate)
Untreated	23	35
Solution 1	6	18
Solution 3	12	20
Solution 1 + Solution 3	0	12

Table B4

Fertiliser Attributes – Spring Barley

Treatment (Applied initially and repeated after a 10 day interval)	Amount of Rooting (0-9) (x 1 Rate)	Amount of Above Ground Tissue (0-9) (x 1 Rate)
Untreated	5.2	5.0
Solution 1	5.3	5.7
Solution 3	6.0	5.7
Solution 1 + Solution 3	6.3	6.0

Table B5

Percent Powdery Mildew - Grape

Treatment (Applied initially and repeated at 10 day intervals)	Days After First Spray					
	+ 16 Days	+ 20 Days	+ 24 Days	+ 28 Days	+ 32 Days	+ 36 Days
Untreated	7.3	9.3	11.3	14.0	14.0	14.7
KP40 @ 0.75 l/ha	5.3	6.7	8.0	11.3	10.7	10.7
KS20 @ 1.0 l/ha	5.3	6.7	8.0	11.3	9.3	11.3
KP40 @ 0.75 l/ha + KS20 @ 1.0 l/ha	1.3	2.7	4.0	5.3	4.7	4.7

Table B6

% Tuber Blight and Yield Attributes - Potato

Treatment (Applied initially and repeated at 10 day intervals)	Percent Tuber Blight (%)	Final Tuber Yield (g)	Final Tuber 'Quality' (Premium Potatoes) (0-9)	Mean Final 'First Grade' Tuber Yield (g)	Relative Final 'First Grade' Tuber Yield (%)
Untreated	6.0	241.7	4.73	114.3	100
KP40 @ 0.75 l/ha	2.7	259.8	4.67	121.3	106
KS20 @ 1.0 l/ha	3.3	255.6	5.07	129.6	113
KP40 @ 0.75 l/ha + KS20 @ 1.0 l/ha	0.7	267.4	5.67	151.6	133

Table B7

Percent Foliar Blight - Potato

Treatment (Applied initially & repeated at 10 day intervals)	Days after First Spray					
	+ 16 Days	+ 20 Days	+ 24 Days	+ 28 Days	+ 32 Days	+ 36 Days
Untreated	5.3	8.7	12.7	12.7	18.7	28.0
KP40 @ 0.75 l/ha	2.7	3.3	6.7	8.7	9.3	13.3
KS20 @ 1.0 l/ha	2.0	4.0	5.3	8.0	10.0	11.3
KP40 @ 0.75 l/ha + KS20 @ 1.0 l/ha	0.7	1.3	1.3	2.0	2.7	4.7

Examples C

Solution 2 = PF723 at 1.0 l/ha every 10 days

Solution 3 = CS8 at 1.0 l/ha every 10 days

Table C1**Percent Powdery Mildew - Grape**

Treatment (Applied initially and repeated at 10 day intervals)	Days After First Spray					
	+ 16 Days	+ 20 Days	+ 24 Days	+ 28 Days	+ 32 Days	+ 36 Days
Untreated	7.3	9.3	11.3	14.0	14.0	14.7
KT47 @ 1.5 l/ha	4.7	6.7	8.0	9.3	9.3	9.3
KS20 @ 1.0 l/ha	5.3	6.7	8.0	11.3	9.3	11.3
KT47 @ 1.5 l/ha + KS20 @ 1.0 l/ha	2.7	5.3	6.7	8.7	7.3	6.7

Table C2**Fertiliser Attributes – Broad Bean**

Treatment (Applied initially and repeated after a 10 day interval)	Amount of Rooting (0-9) (x 1 Rate)	Above Ground Fresh Weight (g) - Mean (x 1 Rate)	Above Ground Dry Weight (g) - Mean (x 1 Rate)
Untreated	5.3	143.8	14.6
Solution 2	5.3	155.7	16.1
Solution 3	5.3	155.0	15.7
Solution 2 + Solution 3	5.7	163.3	16.6

Table C3**Fertiliser Attributes – Peas**

Treatment (Applied initially and repeated after a 10 day interval)	Plant Health 'Greenness' Score (0-9) + 28 Days (x 1 Rate)	Root Fresh Weight (g) - Mean (x 1 Rate)	Root Dry Weight (g) - Mean (x 1 Rate)	Amount of Above Ground Tissue (0-9) at Harvest (x 1 Rate)
Untreated	6.1	124.2	14.1	5.5
Solution 2	6.2	126.3	15.1	5.3
Solution 3	6.2	125.7	15.1	5.7
Solution 2 + Solution 3	6.4	130.7	15.6	6.0

Table C4**Fertiliser Attributes – Carrot**

Treatment (Applied initially and repeated after a 10 day interval)	Root Dry Weight (g) - Mean (x 1 Rate)	Amount of Rooting (0-9) (x 1 Rate)	Amount of Above Ground Tissue (0-9) at Harvest (x 1 Rate)
Untreated	5.5	5.2	5.0
Solution 2	6.1	5.3	5.0
Solution 3	6.2	5.0	5.0
Solution 2 + Solution 3	6.4	5.7	5.7

Table C5

Percent Powdery Mildew – Oilseed Rape

Treatment (Applied initially and repeated after a 10 day interval)	Days After First Spray
	+ 35 Days (x 1 Rate)
Untreated	14
Solution 2	14
Solution 3	14
Solution 2 + Solution 3	8

15 Table C6

Fertiliser Attributes – Sugar Beet

Treatment (Applied initially and repeated after a 10 day interval)	Root Fresh Weight (g) - Mean (x 1 Rate)	Root Dry Weight (g) - Mean (x 1 Rate)
Untreated	109.7	13.3
Solution 2	111.7	13.3
Solution 3	113.7	13.4
Solution 2 + Solution 3	114.7	14.1

Table C7

Fertiliser Attributes - Strawberry

Treatment (Applied initially and repeated after a 10 day interval)	Plant Health 'Greenness' Score (0-9) + 35 Days (x 1 Rate)	Amount of Rooting (0-9) (x 1 Rate)
Untreated	5.0	5.2
Solution 2	6.0	5.7
Solution 3	6.6	5.3
Solution 2 + Solution 3	7.0	6.0

T02T0T"00262850

Examples D

POTATO TRIAL

Table D1

% Tuber Blight, Final Yield and Quality

Treatment (Applied initially and repeated at 10 day intervals)	Percent Tuber Blight (%)	Final Tuber Yield (g)	Relative Final Tuber Yield (%)	Final Tuber 'Quality' (Premium Potatoes) (0-9)	Mean Final 'First Grade' Tuber Yield (g)	Relative Final 'First Grade' Tuber Yield (%)
Untreated	6.0	241.7	100	4.73	114.3	100
KP40 @ 0.75 l/ha	2.7	259.8	107	4.67	121.3	106
KT47 @ 1.5 l/ha	2.7	261.7	108	5.27	137.9	121
KS20 @ 1.0 l/ha	3.3	255.6	106	5.07	129.6	113
KP40 @ 0.75 l/ha + KT47 @ 1.5 l/ha	0.7	271.4	112	5.60	152.0	133
KP40 @ 0.75 l/ha + KS20 @ 1.0 l/ha	0.7	267.4	111	5.67	151.6	133
KT47 @ 1.5 l/ha + KS20 @ 1.0 l/ha	0.7	272.2	113	5.60	152.4	133
KP40 @ 0.75 l/ha + KT47 @ 1.5 l/ha + KS20 @ 1.0 l/ha	0.7	277.0	115	5.80	160.7	141

POTATO TRIAL

Table D2

% Tuber Blight, Yield and Quality Benefits

Treatment (Applied initially and repeated at 10 day intervals)	Percent Tuber Blight (%)	Final Tuber Yield (g)	Final Tuber 'Quality' (Premium Potatoes) (0-9)	Mean Final 'First Grade' Tuber Yield (g)	Relative Final 'First Grade' Tuber Yield (%)
Untreated	14.0	256.1	5.80	148.5	100
KP40 @ 0.75 l/ha	10.0	271.4	5.73	155.5	105
KT47 @ 1.5 l/ha	9.3	279.2	5.80	161.9	109
KS20 @ 1.0 l/ha	9.3	292.0	5.73	167.3	113
KP40 @ 0.75 l/ha + KT47 @ 1.5 l/ha	4.0	280.4	5.80	162.6	109
KP40 @ 0.75 l/ha + KS20 @ 1.0 l/ha	4.7	286.3	5.80	166.1	112
KT47 @ 1.5 l/ha + KS20 @ 1.0 l/ha	3.3	290.0	5.87	170.2	115
KP40 @ 0.75 l/ha + KT47 @ 1.5 l/ha + KS20 @ 1.0 l/ha	1.3	278.0	6.07	168.7	114

Table D3

Spring Barley 1999

Percent Powdery Mildew and Fertiliser Attributes

(KP40 = 40% Potassium Phosphite; CS100 = 10g/litre Salicylamide; CS8 = 20g/l Salicylamide; AT60 = 60% Ammonium Thiosulphate)

Treatment (2 Applications in total - every 15 Days)	Days After First Spray	Plant Health 'Greenness' Score (0-9) + 28 Days	Plant Health 'Greenness' Score (0-9) + 35 Days	Amount of Rooting (0-9) - Mean	Above Ground Fresh Weight (g) - Mean	Above Ground Dry Weight (g) - Mean	Amount of Above Ground Tissue (0-9) - Mean
	+ 20 Days						
Untreated	7	5.4	5.6	5.3	68.8	7.7	5.0
KP40 (0.375 l/ha) + CS100 (1.0 l/ha)	2	5.8	6.0	6.0	72.7	8.4	5.3
KP40 (0.75 l/ha) + CS100 (1.0 l/ha)	2	6.0	6.2	5.3	72.0	8.2	5.0
KP40 (0.375 l/ha) + CS100 (0.5 l/ha)	2	5.8	5.4	6.0	75.3	8.5	5.7
KP40 (0.75 l/ha) + CS100 (0.5 l/ha)	4	5.6	5.8	6.3	68.7	7.5	5.3
KP40 (0.375 l/ha) + CS100 (1.0 l/ha) + AT60 (10.0 l/ha)	2	5.4	6.0	6.7	69.7	8.0	6.0
KP40 (0.375 l/ha) + CS100 (0.5 l/ha) + AT60 (10.0 l/ha)	2	5.6	6.2	6.0	68.7	7.9	5.7
KP40 (0.375 l/ha) + CS100 (0.5 l/ha) + AT60 (6.0 l/ha)	4	5.8	5.6	6.0	62.0	7.4	5.0
KP40 (0.375 l/ha) + CS8 (0.05 l/ha) + AT60 (10.0 l/ha)	0	5.2	5.4	6.0	69.0	7.8	5.3
KP40 (0.375 l/ha) + CS8 (0.05 l/ha) + AT60 (6.0 l/ha)	2	6.0	5.6	6.7	67.7	7.5	5.0
KP40 (0.375 l/ha) + CS8 (0.25 l/ha) + AT60 (10.0 l/ha)	0	5.4	5.0	5.7	69.7	7.4	5.0
KP40 (0.375 l/ha) + CS8 (0.25 l/ha) + AT60 (6.0 l/ha)	0	6.0	5.3	6.0	69.3	7.5	5.0

Spring Barley

(KP40 = 40% Potassium Phosphite; CS100 = 10g/litre Salicylamide; CS8 = 20g/l Salicylamide; AT60 = 60% Ammonium Thiosulphate)

Treatment (2 Applications in total - every 15 Days)	Amount of Rooting (0-9) - Mean	Above Ground Fresh Weight (g) - Mean	Above Ground Dry Weight (g) - Mean	Amount of Above Ground Tissue (0-9) - Mean
Untreated	5.2	73.0	9.1	5.0
KP40 (0.375 l/ha) + CS100 (1.0 l/ha)	6.0	83.0	9.3	5.3
KP40 (0.75 l/ha) + CS100 (1.0 l/ha)	5.7	82.0	9.7	5.3
KP40 (0.375 l/ha) + CS100 (0.5 l/ha)	5.0	77.7	8.6	6.0
KP40 (0.75 l/ha) + CS100 (0.5 l/ha)	6.0	76.7	8.7	5.0
KP40 (0.375 l/ha) + CS100 (1.0 l/ha) + AT60 (10.0 l/ha)	6.0	67.3	7.8	5.0
KP40 (0.375 l/ha) + CS100 (0.5 l/ha) + AT60 (10.0 l/ha)	6.0	68.3	8.1	5.0
KP40 (0.375 l/ha) + CS100 (0.5 l/ha) + AT60 (6.0 l/ha)	6.0	78.3	9.1	5.7
KP40 (0.375 l/ha) + CS8 (0.05 l/ha) + AT60 (10.0 l/ha)	5.7	76.0	8.8	5.7
KP40 (0.375 l/ha) + CS8 (0.05 l/ha) + AT60 (6.0 l/ha)	6.0	78.3	9.0	5.7
KP40 (0.375 l/ha) + CS8 (0.25 l/ha) + AT60 (10.0 l/ha)	5.0	71.7	8.5	5.3
KP40 (0.375 l/ha) + CS8 (0.25 l/ha) + AT60 (6.0 l/ha)	5.7	72.0	8.1	5.0

LETTUCE TRIAL

Table D5

Fresh Weight Yield and Quality Benefits

(KP40 = 40% Potassium Phosphate; PF723 = 55% Ammonium Thiosulphate; KS20 = 20 gms/litre Potassium Salicylate)

Treatment (Applied Initially and repeated at 10 day intervals)	Final Fresh Weight Yield (g)	Quality of Final Fresh Weight Yield (0-9)	Final Fresh Weight 'Quality' Yield (g)	Relative Fresh Weight 'Quality' Yield (%)	Median Final Fresh Weight Yield (g)	Median Quality of Final Fresh Weight Yield (0-9)	Median Final Fresh Weight 'Quality' Yield (g)	Relative Median Fresh Weight 'Quality' Yield (%)
Untreated	81.3	5.33	43.4	100	80.8	5.27	42.5	100
KP40 @ 0.75 l/ha	85.0	5.40	45.7	105	87.2	5.36	46.7	110
PF723 @ 1.0 l/ha	87.7	5.40	47.4	108	88.6	5.46	47.3	111
KS20 @ 1.0 l/ha	83.6	5.67	47.4	108	82.6	5.64	46.6	110
KP40 @ 0.75 l/ha + PF723 @ 1.0 l/ha	89.8	5.63	50.6	117	89.8	5.64	50.6	119
KP40 @ 0.75 l/ha + KS20 @ 1.0 l/ha	90.9	5.53	50.3	116	90.6	5.64	51.1	120
PF723 @ 1.0 l/ha + KS20 @ 1.0 l/ha	86.5	5.67	49.0	113	85.4	5.64	48.2	113
KP40 @ 0.75 l/ha + PF723 @ 1.0 l/ha + KS20 @ 1.0 l/ha	87.9	5.67	49.8	115	88.4	5.64	49.9	117

Table E1

5 Benefits of Ammonium Thiosulphate (ATS) and Potassium Salicylate (KS) with
Chlormequat (CCC) on Spring Barley

Treatment (Treatments applied at 3 leaves stage)	Powdery Mildew (%) at + 15 days	Amount of Rooting (0-9) Mean of 10 plants	Above Ground Fresh Weight (g) Total of 10 plants	Above Ground Dry Weight (g) Total of 10 plants
Untreated	25	5.8	30.5	3.4
CCC (1.25 l/ha)	15	5.5	34.0	3.7
CCC (1.25 l/ha) + ATS (1.25 l/ha)	13	6.0	31.5	3.6
CCC (1.25 l/ha) + KS (20 g/ha)	18	6.0	31.3	3.5
CCC (1.25 l/ha) + ATS (1.25 l/ha) + KS (20 g/ha)	8	6.5	36.0	4.0

Table E2

Benefits of Ammonium Thiosulphate (ATS) and Potassium Salicylate (KS) with
Chlormequat (CCC) on Spring Barley

Treatment (Treatments applied at 3 leaves stage (T1) and prior to start of stem extension (T2))	Powdery Mildew (%) at + 12 days	Powdery Mildew (%) at + 15 days	Powdery Mildew (%) at + 18 days	Number of Tillers Initiated per Plant (mean of 10 plants)
Untreated	20	25	38	3.0
CCC (0.8 l/ha at T1 & T2)	8	8	10	3.5
CCC (0.8 l/ha at T1 & T2) + ATS (0.8 l/ha at T1 & T2)	5	5	8	3.5
CCC (0.8 l/ha at T1 & T2) + KS (20 g/ha at T1 & T2)	8	8	13	3.3
CCC (0.8 l/ha at T1 & T2) + ATS (0.8 l/ha at T1 & T2) + KS (20 g/ha at T1 & T2)	0	3	5	4.3

0939900-101201
T02T0T-00868860

The following show non-limiting examples of formulated compositions in accordance with the present invention

FOLIAR FERTILISER ONE

INGREDIENTS	Specific Gravity	Kilogram per batch	percentage w/w	Volume per batch	gram/litre of ingredient	gram/litre of active
Water	1.000	150.0000	7.50000	150.0000	96.2564	96.2564
Wetting agent	1.000	10.0000	0.50000	10.0000	6.4171	6.4171
Salicylic Acid	1.000	10.0000	0.50000	10.0000	6.4171	6.4171
Potassium hydroxide (20% w/w)	1.200	130.0000	6.50000	108.3333	83.4222	83.4222
Copper EDTA chelate (14.3 % Cu w/w;	1.200	30.0000	1.50000	25.0000	19.2513	19.2513
Iron EDTA chelate (13.2 %Fe w/w)	1.200	30.0000	1.50000	25.0000	19.2513	19.2513
Pot. phosphites (42%w/w)	1.342	1,000.0000	50.00000	745.1565	641.7092	269.5179
Ammonium thiosulphate (60% w/w)	1.320	640.0000	32.00000	484.8485	410.6939	246.4163
Totals		2,000.0000	100.00000	1,558.3383	1,283.4184	746.9495
N			3.84 % w/w			49 g/litre w/v
P			4.68 % w/w			60 g/litre w/v
P as P ₂ O ₅ **			10.64 % w/w			137 g/litre w/v
K			9.91 % w/w			125 g/litre w/v
K as K ₂ O			11.93 % w/w			150 g/litre w/v
S			8.32 % w/w			107 g/litre w/v
S as SO ₃			20.80 % w/w			267 g/litre w/v
** theoretically						

FOLIAR FERTILISER TWO

INGREDIENTS	Specific Gravity	Kilogram per batch	percentage w/w	Volume per batch	gram/litre of ingredient	gram/litre of active
Water	1.000	150.0000	7.50000	150.0000	96.2564	96.2564
Wetting agent	1.000	10.0000	0.50000	10.0000	6.4171	6.4171
Zinc EDTA chelate (15.7% Zn w/w)	1.000	30.0000	1.50000	30.0000	19.2513	19.2513
Copper EDTA chelate (14.3% Cu w/w)	1.000	30.0000	1.50000	30.0000	19.2513	19.2513
Iron EDTA chelate (13.2% Fe w/w)	1.000	30.0000	1.50000	30.0000	19.2513	19.2513
Pot. phosphites (42%w/w)	1.342	750.0000	37.50000	558.8674	481.2819	202.1384
Ammonium thiosulphate (60% w/w)	1.320	1,000.0000	50.00000	757.5758	641.7092	385.0255
Totals		2,000.0000	100.00000	1,566.4431	1,283.4184	747.5912
N			6.00 % w/w			77 g/litre w/v
P			3.51 % w/w			45 g/litre w/v
P as P ₂ O ₅ **			7.98 % w/w			102 g/litre w/v
K			6.96 % w/w			88 g/litre w/v
K as K ₂ O			8.39 % w/w			105 g/litre w/v
S			13.00 % w/w			167 g/litre w/v
S as SO ₃			32.50 % w/w			417 g/litre w/v
** theoretically						

The above Examples show that the compositions of the present invention show the desired fertilisation and antifungal effects.

CLAIMS

1. A fertiliser composition comprising at least one phosphonate and at least one thiosulphate.
2. A fertiliser composition according to claim 1 further comprising at least one salicylic acid, homologue, derivative, or salt thereof.
3. A fertiliser composition comprising at least one phosphonate and at least one salicylic acid, homologue, derivative, or salt thereof.
4. A fertiliser composition according to claim 3 further comprising at least one thiosulphate.
5. A fertiliser composition comprising at least one thiosulphate and at least one salicylic acid, homologue, derivative, or salt thereof.
6. A fertiliser composition according to claim 5 further comprising at least one phosphonate.
7. A fertiliser composition according to any one of claims 1-4 and 6 wherein the phosphonate is ammonium, sodium or potassium phosphonate or a mixture thereof.
8. A fertiliser composition according to any one of claims 1 and 4-7 wherein the thiosulphate is ammonium, sodium or potassium thiosulphate or a mixture thereof.

37

9. A fertiliser composition according to any one of claims 2-8 wherein the derivative of salicylic acid is salicylamide or a salt thereof.

5 10. A fertiliser composition according to any one of claims 2-8 wherein the homologue of salicylic acid is benzoic acid or a salt or derivative thereof.

11. A fertiliser composition according to any of claims 2-9 wherein the salt of salicylic acid, its homologue or derivative is an organic or inorganic salt.

10 12. A fertiliser composition according to claim 11 wherein the salt is a sodium or potassium salt or mixtures thereof.

13. A fertiliser composition according to any preceding claim in the form of a concentrate.

15 14. A fertiliser composition according to any one of claims 1-12 in the form of an aqueous solution.

20 15. A fertiliser composition according to claim 14 comprising 150 g/l phosphonate, 275 g/l thiosulphate and/or 10 g/l salicylamide.

25 16. A fertiliser composition according to claim 15 wherein the phosphonate comprises 75 g/l mono-potassium phosphonate and 75 g/l di-potassium phosphonate.

17. A fertiliser composition according to any preceding claim further comprising a plant growth regulator.

30 18. A fertiliser composition according to claim 17 wherein the plant growth regulator is chlormequat.

19. A method for fertilising a plant comprising applying a fertiliser composition according to any preceding claim to the plant or its environs.

5 20. A method according to claim 19 wherein the phosphonate is applied at 150 g/ha to 2 kg/ha.

21. A method according to claim 19 or claim 20 wherein the thiosulphate is applied at 250 g/ha to 6 kg/ha.

10 22. A method according to any one of claims 19 to 21 wherein at least one salicylic acid, a homologue, derivative, or salt thereof is applied at 1 g/ha to 100 g/ha.

15 23. Use of a fertiliser composition according to any one of claims 1-18 to stimulate plant growth.

20 24. Use of a fertiliser composition according to any one of claims 1-18 to control parasitic fungi.

PATENT (US)

Attorney Docket No. DYOUP0219US

COMBINED DECLARATION AND POWER OF ATTORNEY
(ORIGINAL, DESIGN, NATIONAL STAGE OF PCT)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

Title: FERTILISER

the specification of which

(a) is attached hereto.

(b) x was filed on July 20, 2001 as Serial No. 0 / or X Express Mail No. EF232849690US, as Serial No. not yet known, and was amended on (if applicable).

(c) x was described and claimed in PCT International Application No. PCT/GB00/00367 filed on February 7, 2000 and amended under PCT Article 34 on April 27, 2001 (if any).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations §1.56(a).

PRIORITY CLAIM

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed.

(d) no such application have been filed.

(e) x such applications have been filed as follows.

EARLIEST FOREIGN APPLICATION(S), IF ANY FILED WITHIN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

COUNTRY	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119
<u>United Kingdom</u>	<u>9902665.0</u>	<u>05/02/99</u>	<u>x</u> Yes <u> </u> No

ALL FOREIGN APPLICATION(S), IF ANY FILED MORE THAN 12 MONTHS
(6 MONTHS FOR DESIGN) PRIOR TO THIS U.S. APPLICATION

POWER OF ATTORNEY

As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. (List name and registration number)

Don W. Bulson, Reg. No. 28,192; Neil A. DuChez, Reg. No. 26,725; Mark D. Saralino, Reg. No. 34,243

FOOTNOTES

The undersigned to this declaration and power of attorney hereby authorizes the U.S. attorney(s) named herein to accept and follow instruction from

Name(s) of authorized representation(s) D. Young & Co.

Address 21 New Fetter Lane, London EC4A 1DA

as to any actions to be taken in the Patent and Trademark Office regarding this application without direct communication between the U.S. attorney(s) and the undersigned. In the event of a change in the person(s) from whom instructions may be taken, the U.S. attorney(s) will be so notified by the undersigned.

Send Correspondence To

Neil A. DuChez
RENNER, OTTO, BOISSELLE & SKLAR, LLP
1621 Euclid Avenue, 19th Floor
Cleveland, Ohio 44115

Direct Telephone Calls To:
(name and telephone number)

Neil A. DuChez
(216) 621-1113

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with knowledge that willful false statements and the like are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issued therein.

Full name of sole or first inventor Richard Henry Williams

Inventor's signature [Signature]
Date 7/8/01 Country of Citizenship Great Britain

Residence Essex CO4 3FH, Great Britain

Post Office Address 15 Chaffinch Gardens, Colchester
Essex CO4 3FH, Great Britain

Full name of second inventor, if any Peter Harding

Inventor's signature [Signature]
Date 7 AUGUST 2001 Country of Citizenship Great Britain

Residence Hampshire SO18 5QR, Great Britain

Post Office Address 6 Cutbush Lane, Bitterne, Southampton
Hampshire SO18 5QR, Great Britain

Full name of third inventor, if any _____

Inventor's signature _____

Date _____ Country of Citizenship _____

Residence _____

Post Office Address _____

**CHECK FOR ANY OF THE FOLLOWING ADDED PAGE(S) WHICH
FORM A PART OF THIS DECLARATION**

____ Signature for fourth and subsequent joint inventors. Number of pages added ____.

____ Added page to combined declaration and power of attorney for divisional, continuation, or continuation-in-part (CIP) application.

X This declaration ends with this page.